

PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Search PubMed



for



Limits

Preview/Index

History

Clipboard

Data

About Entrez

Display

Abstract



Sort



Save

Text

Clip Add

Order

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: Neth J Med 1993 Aug;43(1-2):83-90

Related Articles, Books, Link

The role of modification of lipoproteins and of the immune system in early atherogenesis.

Mol MJ, Demacker PN, Stalenhoef AF.

Department of Medicine, University Hospital, Nijmegen, Netherlands.

Not only the plasma cholesterol level, but also postsecretory modification of lipoproteins appear to be of influence in atherogenesis. Evidence that several forms of modification, especially oxidation, occur in vivo is rapid accumulating, although their clinical relevance remains uncertain.

Modification of lipoproteins has been demonstrated in persons with such well-known risk factors of premature atherosclerosis as smoking, diabetes mellitus and hyperlipidaemia. Because there is a relation between the amount of natural antioxidants in the plasma and the risk of atherosclerosis, and because exogenous antioxidants appear to retard atherosclerosis without influencing the plasma cholesterol level, antioxidants may prove to be of use in the prevention of atherosclerosis. There are strong indications for a role of the immune system in atherogenesis. Modified lipoproteins are highly immunogenic and stimulate immunocompetent cells to secrete vasoactive factors and cytokines. From animal studies it appears that pro- and antioxidative conditions can modulate these processes. It is concluded that additional research on the relation between lipoprotein modification and the immune system, and on the possible beneficial effects of antioxidants in atherogenesis is warranted, not only to elucidate further the mechanism of atherosclerosis, but also to develop new approaches to the prevention of atherosclerosis.

Publication Types:

- Review
- Review Literature

PMID: 8232700 [PubMed - indexed for MEDLINE]

Display

Abstract



Sort


Save

Text

Clip Add

Order

Write to the Help Desk
NCBI | NLM | NIH



PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Search PubMed for

Go

Clear

Limits

Preview/Index

History

Clipboard

Data

About Entrez

Display

Abstract

Sort

Save

Text

Clip/Add

Order

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: Ann Biol Clin (Paris) 1992;50(4):213-27

Related Articles, Books, Link

[Modified LDL and atherosclerosis. Nature of modifications. Physicochemical and biological properties]

[Article in French]

Pech MA, Myara I, Védie B, Moatti N.

Laboratoire de biochimie appliquee, faculte des sciences pharmaceutique et biologiques, Chatenay-Malabry, France.

The role of plasma LDL in atherogenesis is now well established. Cholesteryl ester accumulation within macrophages leads to foam cell formation, an early atherosclerotic process. In vitro, foam cell formation scarcely ever occurs in the presence of native LDL. Modification of these lipoproteins is necessary for their binding to macrophage scavenger receptors. In vitro modifications reported have involved chemical reactions, physical mechanisms, enzymatic reactions, cellular interactions and association with macromolecules. In vivo, they can occur by glycation or desialylation, smoking or by hemodynamic interactions. Alterations of the physicochemical properties of LDL are induced by oxidation and include an increase in their density and their net electronegative charge, changes in lecithin composition, polyunsaturated fatty acid peroxidation of lipids and apolipoprotein B100 degradation. Apolipoprotein B100 fragmentation leads to an impairment of uptake through LDL receptors, while uptake through macrophage scavenger receptors is enhanced. Modified LDL have also other particularities such as cytotoxicity, chemotactism for circulating monocytes, inhibition of resident macrophage mobility, vasoconstriction, perturbation of the arachidonic acid cascade, involvement in haemostasis and immune mechanisms. Hypotheses concerning the role of modified LDL, in particular oxidized LDL, in atherogenesis open new therapeutic prospects.

Publication Types:

- Review
- Review, Academic

PMID: 1456499 [PubMed - indexed for MEDLINE]

Display

Abstract

Sort

Save

Text

Clip/Add

Order